

Suppression of isoscalar pairing

G.F. Bertsch and Simone Baroni

Department of Physics and Institute for Nuclear Theory, University of Washington, Seattle, WA 98195

The short-range nuclear attraction is stronger in the isoscalar channel than in the isovector channel, as evidenced by the existence of the deuteron and not the dineutron. Nevertheless, apart from light $N = Z$ nuclei, pairing is only seen in the isovector channel. This is explained by the effect of the strong spin-orbit splitting on the single-particle energies. A semiquantitative argument is presented treating the high- j orbitals at the Fermi surface as plane waves on a two-dimensional sheet.

Calculations with realistic interactions predict that in nuclear matter pairing should be very strong in the $(S, T) = (1, 0)$ channel [1]. Nevertheless, pairing in nuclear structure is only visible in the $J = 0$ isovector channel, except for a few light odd-odd $N = Z$ nuclei, for which the ground state has $J = 1$. The reason is that the spin-orbit field affects the pairing in the isoscalar channel more strongly than in the isovector channel. The effect is implicit in shell-model calculations such as those of ref. [3], dealing with the nonappearance of $T = 0$ pairing. Still, it is of interest to see how the effect comes about in a qualitative, analytic treatment of the interaction in jj coupled orbitals. We first note that the high- j orbitals are localized at the nuclear surface, and their local momentum is mainly tangential to the surface. In the limit of a very large nucleus, one can treat the interactions as those of particles localized on a 2-dimensional plane tangent to the surface. On this plane, the Fermi surface is a circle and we need only to investigate the interactions on the circle.

We will analyze the momentum and spin of the particles in the vicinity of the point $\vec{r} = (0, 0, R)$ where R is the radius of the nucleus. The surface of the sphere is mapped onto a plane perpendicular to the z axis as shown in Fig. 1. In this plane, the orbitals can be expressed in terms of their linear momentum (k_x, k_y) and the spin projection s_z as the kets $|k_x, k_y, s_z\rangle$. In the presence of the spin-orbit field, the eigenfunctions will have the spin oriented in the xy plane and perpendicular to the momentum $\vec{k}$ in the xy -plane. For the tangent plane shown in Fig. 1, the preferred orientation corresponding to states with $j = \ell + 1/2$ is $\hat{z} \times \hat{k}$. Thus, the preferred state with k along the x -axis has spin oriented along the y -axis,

$$|x \uparrow_y\rangle = \frac{e^{i\pi/4}}{\sqrt{2}}|k, 0, \uparrow_z\rangle + \frac{e^{-i\pi/4}}{\sqrt{2}}|k, 0, \downarrow_z\rangle. \quad (1)$$

In the general case, when $\vec{k}$ is oriented at an angle ϕ with respect to the x -axis, the preferred state is given by

$$\begin{aligned} |\phi \uparrow_p\rangle &= \mathcal{R}_\phi |x \uparrow_y\rangle \\ &= \frac{e^{i(\pi/4 + \phi/2)}}{\sqrt{2}} |k \cos \phi, k \sin \phi, \uparrow_z\rangle + \\ &\quad \frac{e^{-i(\pi/4 + \phi/2)}}{\sqrt{2}} |k \cos \phi, k \sin \phi, \downarrow_z\rangle \end{aligned} \quad (2)$$

where the subscript on $\uparrow_p$ stands for the *preferred* axis of spin orientation. This is shown in Fig. 1.




FIG. 1: The geometry of high- j orbital state $|\phi, \uparrow_p\rangle$ projected onto a tangent plane. The directions of the spin $\vec{s}$ and linear momentum $\vec{k}$ correspond to a spherical orbital with $j = \ell + 1/2$.

To treat the pairing interaction, we consider the set of 2-particle states defined by the product of one-particle states having momenta k and $-k$. The state with preferred spin orientation for both particles is

$$|2p, \phi, \uparrow_p \uparrow_p\rangle = |\phi \uparrow_p\rangle |(\phi + \pi) \uparrow_p\rangle \quad (3)$$

as shown in Fig. 2. We will later need to transform the spins to a common frame using Eqs. (1) and (2). In the usual z -oriented basis, the spin wave function of the preferred 2-particle state is

$$\begin{aligned} |2p, \phi, \uparrow_p \uparrow_p\rangle &= \\ &= \frac{1}{2} (e^{i\phi} |\uparrow_z \uparrow_z\rangle + e^{-i\phi} |\downarrow_z \downarrow_z\rangle \\ &\quad + e^{-i\pi/2} |\uparrow_z \downarrow_z\rangle + e^{i\pi/2} |\downarrow_z \uparrow_z\rangle) \\ &= \frac{1}{2} (e^{i\phi} |1, 1\rangle + e^{-i\phi} |1, -1\rangle - i\sqrt{2} |0, 0\rangle). \end{aligned} \quad (4)$$

In the last line, we expressed the spin wave function in the coupled form $|S, S_z\rangle$.

The interaction will mix wave functions having different orientations ϕ , and the mixed state for spin s_1, s_2 will have the form

$$N \int_0^{2\pi} d\phi f(\phi) |2p, \phi, s_1, s_2\rangle \quad (5)$$




FIG. 2: Visualization, of the state $|2p, \phi, \uparrow_p \uparrow_p\rangle$ in a two-dimensional sheet, corresponding to two particles on the nuclear surface, scattering in time reversal states, with spin orientations dictated by the spin-orbit interaction.

where $f(\phi)$ is an amplitude and N is a normalization factor. Due to the rotational symmetry in the xy plane, we can take the amplitudes to have the form $f(\phi) = e^{iM\phi}$. Note that the wave function has a definite particle exchange symmetry because the oriented function transforms as $P_{12}|2p, \phi, s_1 s_2\rangle = -|2p, \phi + \pi, s_1 s_2\rangle$. Thus, the even- M mixed states are antisymmetric (ie. $T = 1$) while the odd- M states are symmetric ($T = 0$). We determine the norm N by defining the overlap matrix element,

$$\langle 2p, \phi', s_1, s_2 | 2p, \phi, s_3, s_4 \rangle = \delta(\phi - \phi') \delta_{s_1, s_3} \delta_{s_2, s_4}. \quad (6)$$

Then the normalized wave functions for the interacting states are

$$|2p, M, s_1, s_2\rangle = \frac{1}{\sqrt{2\pi}} \int_0^{2\pi} d\phi e^{iM\phi} |2p, \phi, s_1, s_2\rangle. \quad (7)$$

We are now ready to calculate matrix elements. For a warm-up exercise, we start with a spin-independent interaction

$$v = \frac{v_W}{2\pi} \delta^{(2)}(\vec{r}_1 - \vec{r}_2) \quad (8)$$

and go back to the original spin representation, quantizing along the z -axis. The matrix element is

$$\langle 2p, \phi', s_{1z}, s_{2z} | v | 2p, \phi, s_{3z}, s_{4z} \rangle = \frac{v_W}{2\pi} \delta_{s_{1z}, s_{3z}} \delta_{s_{2z}, s_{4z}} \quad (9)$$

where we have set the area corresponding to the integration over the center-of-mass to one. The matrix element is independent of ϕ and ϕ' , which leads to a uniform mixing of ϕ values in the paired state, ie. $M = 0$ in the wave function Eq. (7). The expectation value of the interaction energy in the $M = 0$ state is given by

$$\langle 2p, M = 0, s_{1z}, s_{2z} | v | 2p, M = 0, s_{1z}, s_{2z} \rangle = v_W \quad (10)$$

It is easy to include a spin dependence in the interaction in the s_z representation. We write the spin-dependent interaction as

$$v = \frac{1}{2\pi} (v_0 P(0) + v_1 P(1)) \quad (11)$$

where $P(S)$ projects on the total spin S . If our space includes all spin projections, i.e. the states $|2p, M = 0, s_1, s_2\rangle$ with the four orientations $s_1, s_2 = \pm 1$, we can recouple the spin states into a singlet and a triplet with 3 substates. In that representation, the interaction energy including the spin projector is v_0 for the singlet state and v_1 for each of the three triplet states.

Now we come to the main question, how are these interaction energies reduced when the space is restricted to the preferred spin orientation? For this we need the matrix element of the interaction in the ϕ -representation, Eq. (4),

$$\begin{aligned} \langle 2p, \phi', \uparrow_p \uparrow_p | v | 2p, \phi, \uparrow_p \uparrow_p \rangle &= \\ &= \frac{1}{4} \left(e^{i(\phi - \phi')} \langle 1, 1 | v | 1, 1 \rangle \right. \\ &\quad \left. + e^{-i(\phi - \phi')} \langle 1, -1 | v | 1, -1 \rangle + 2 \langle 0, 0 | v | 0, 0 \rangle \right) \\ &= \frac{1}{8\pi} \left(e^{i(\phi - \phi')} v_1 + e^{-i(\phi - \phi')} v_1 + 2v_0 \right). \end{aligned} \quad (12)$$

To find the eigenstates, we apply the interaction to states of the form Eq. (7). It is obvious that the three terms in Eq. (11) select different M values, with the v_1 interaction surviving for $M = \pm 1$ and the v_0 interaction surviving for $M = 0$. The expectation values of the interaction energy are

$$\langle 2p, M, \uparrow_p \uparrow_p | v | 2p, M, \uparrow_p \uparrow_p \rangle = \quad (13)$$

$$\frac{v_0}{2} \text{ for } M = 0,$$

$$\frac{v_1}{4} \text{ for } M = \pm 1.$$

Thus, the reduction from the interaction strength in the full spin space is a factor of two for the $S = 0$ but a factor 4 for the $S = 1$. As long as the intrinsic strength of the spin-triplet interaction is not greater than twice the strength of the spin-singlet interaction, the ordinary $J = 0$ pairing will be favored within a single j -shell (for j large). The factor of 2 suppression of the singlet pairing can be seen directly from the values of the appropriate LS - jj recoupling coefficient. For the shell with $j = \ell + 1/2$, the relevant recoupling coefficient is

$$\langle (jj)^{J=0} | (\ell\ell)^{L=0}, S=0 \rangle = \sqrt{\frac{2j+1}{4j}} \rightarrow \sqrt{\frac{1}{2}}. \quad (14)$$

The situation is somewhat more complicated for the triplet pairing because both $L = 0$ and $L = 2$ couplings

contribute to the single j -shell interaction. When both couplings are taken into account in a contact interaction, one finds that the suppression is indeed a factor four in the limit of large j [2].

-
- [1] M. Baldo, U. Lombardo, and P. Schuck, Phys. Rev. C **52** 975 (1995).
 [2] G.F. Bertsch, A.O. Macchiavelli, and A. Schwenk, unpublished.
 [3] A. Poves, G. Martinez-Pinedo, Phys. Letter B430 203 (1998).